



King's Research Portal

DOI:

[10.3233/JPD-2011-11037](https://doi.org/10.3233/JPD-2011-11037)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Martinez-Martin, P., Reddy, P., Antonini, A., Henriksen, T., Katzenschlager, R., Odin, P., Todorova, A., Naidu, Y., Tluk, S., Chandiramani, C., Martin, A., & Ray Chaudhuri, K. (2011). Chronic subcutaneous infusion therapy with apomorphine in advanced Parkinson's disease compared to conventional therapy: a real life study of non motor effect. *Journal of parkinsons disease*, 1(2), 197-203. <https://doi.org/10.3233/JPD-2011-11037>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Chronic Subcutaneous Infusion Therapy with Apomorphine in Advanced Parkinson's Disease Compared to Conventional Therapy: A Real Life Study of Non Motor Effect

Pablo Martinez-Martin^a, Prashanth Reddy^f, Angelo Antonini^b, Tove Henriksen^c, Regina Katzenschlager^d, Per Odin^e, Antonia Todorova^f, Yogini Naidu^g, Susanne Tluk^g, Chandni Chandiraman^f, Anne Martin^f and Kallol Ray Chaudhuri^{f,g,*}

^a*Alzheimer Disease Research Unit and CIBERNED, Carlos III Institute of Health, Alzheimer Center Reina Sofia Foundation, Madrid, Spain*

^b*Department for Parkinson's disease, IRCCS San Camillo, Venice, Italy*

^c*Department of Neurology, Bispebjerg Hospital, Copenhagen, Denmark*

^d*Department of Neurology, SMZ-Ost Donauspital, Vienna, Austria*

^e*Department of Neurology, Central Hospital, Bremerhaven, Germany*

^f*Department of Neurology, NPF Centre of Excellence, King's College Hospital, MRC Centre for Neurodegeneration Research and Biomedical Research Centre, Kings College, London, United Kingdom*

^g*University Hospital Lewisham, London, UK*

Abstract. *Background:* Apomorphine infusion therapy remains under-used and there are no comparative studies of motor and non-motor effects of apomorphine infusion.

Methods: In this paper we report preliminary results from an ongoing clinical observational “real life” surveillance-based study focused on effects of this therapy on non-motor symptoms and health-related quality of life in a group of patients on apomorphine.

Results: Apomorphine infusion led to highly significant improvements in UPDRS 3 ($p=0.0003$), UPDRS 4 ($p=0.0003$), PDQ-8 (Parkinson's disease questionnaire, $p=0.001$) and NMSS total (non motor symptoms scale, $p=0.0003$). Furthermore, apomorphine was tolerated in patients with visual hallucinations, illusions and paranoid ideations while significant improvement in specific non-motor symptoms such as hyperhidrosis, nocturia, urgency of micturition, and fatigue was recorded. Levodopa equivalent dose decreased significantly (1077.81 ± 446.26 to 458.75 ± 282.29 , $p < 0.0001$) and a large effect size of intervention was noted. In an untreated group no such improvement was noted. The number needed to treat (NNT) for improvement >1 SEM in the Apo group was calculated and was lower than 2 for >1 SEM improvement of UPDRS 3, NMSS, and PDQ-8 total scores.

Conclusions: This pilot observational study suggests that non-motor effects are evident with apomorphine therapy and patients suitable for apomorphine deteriorate in the absence of therapy.

INTRODUCTION

Subcutaneous administration of apomorphine (Apo) has been used as treatment for advanced Parkinson's disease (PD) with refractory on-off periods and dyskinesias and is nationally recommended in many

*Correspondence to: Prof. K. Ray Chaudhuri, Department of Neurology, King's College Hospital, Denmark Hill, SE5 9RS London, UK. E-mail: chaudhuri@hotmail.com.

countries [1–10]. We have reported the beneficial effect of apomorphine infusion on aspects of non-motor symptoms (NMS) of PD as assessed by the validated PD NMS scale (NMSS) [11–14]. There are no studies comparing non-motor effects of apomorphine infusion in two groups of PD patients deemed suitable for apomorphine therapy yet where one group is treated by conventional (oral and patch therapy) treatment only because of local funding issues.

PATIENTS AND METHODS

Between 2008–2009, 17 PD patients (all satisfying the UK PD brain Bank criteria) [15] with severe dyskinesias (as rated by patients, carers and clinician) and on-off fluctuations unresponsive to changes in existing therapy were started on apomorphine infusion (12–16 hrs/day) and followed up in centres across Europe. All had baseline assessment (“7 to 10 days pre-treatment” of motor function (Unified Parkinson’s Disease Rating Scale, UPDRS 3 and 4) [16], Hoehn and Yahr staging (HY) [17], Non-Motor Symptoms Scale (NMSS) [13, 14] and health-related quality of life (HRQoL) (Parkinson’s disease questionnaire, PDQ-8) [18]. Follow-up assessments were carried out in a manner similar to that recently reported [8]. For analysis, baseline data was compared to the last available follow-up visit of each patient at “on” stage.

Seventeen PD patients were selected for comparison (C) as these patients were not funded for apomorphine. These patients also had severe dyskinesias and on-off fluctuations unresponsive to further modifications in their existing therapy. These patients were followed up while on best conventional therapy using the same clinical protocol. As this was a clinically dictated study, matching for C and Apo cases from individual centres was not possible.

As this was a clinical observational surveillance audit based study, with a licensed drug for the approved indication specific ethical approval was not required, as confirmed by written statements from the Bispebjerg Hospital, Copenhagen ethics committee and ethics committee in Donaushpital, Vienna. Use of NMSS methodology was approved for use by the ethics committee of the University Hospital of Lewisham, Kings College Hospital, Madrid, and Bremerhaven.

Statistical analysis

Chi-squared and Fisher’s exact tests were used for comparison of proportions, while non-parametric tests were applied for ordinal and continuous vari-

ables. Multiple comparisons were corrected by the Benjamini-Hochberg method [19] and for the association of changes Spearman rank correlation coefficient was used [20]. For each measure, the relative change [$RC = \text{mean} (\text{test}_{T2} - \text{test}_{T1}) \times 100 / \text{mean test}_{T1}$], the effect size [$ES = \text{mean} (\text{test}_{T2} - \text{test}_{T1}) / SD \text{ test}_{T1}$] and the standardized response mean [$SRM = \text{mean} (\text{test}_{T2} - \text{test}_{T1}) / SD (\text{test}_{T2} - \text{test}_{T1})$] were calculated to determine the magnitude of change. For both effect size indices, ES and SRM, the standard values are: 0.20–0.49, Small effect; 0.50–0.79, Moderate effect; and ≥ 0.80 , Large effect [21–23]. The number needed to treat (NNT) for improvement >1 SEM in the Apo group was calculated [24–27].

RESULTS

Seventeen Apo (mean age 59.5 ± 11.7 yrs, disease duration 12.05 ± 4 yrs, median HY score 4) were started on apomorphine infusion using a standardised local initiation protocol and the data compared to 17 C patients (mean age 66.4 ± 7.0 yrs, disease duration 13.23 ± 4.7 yrs, median HY score 3.9) over a period of 12.5 ± 11.5 months. At baseline, there were no significant differences between Apo and C with respect to age, gender and PD duration and both groups were on comparable levodopa and dopamine agonist treatments and had resistant on-off fluctuations and dyskinesias. The levodopa equivalent (LDE) doses in both groups were similar between Apo (1077.81 ± 446.26 mg) and C (1028.18 ± 388.76 mg) at baseline. At baseline, the UPDRS 3, NMSS-Total and PDQ-8 showed significant differences between C and Apo group, the C group being lower in scores (indicative of better state). During follow-up, the C group showed worsening in PDQ-8 (Table 1) while Apo showed highly significant improvements in UPDRS 3 ($p = 0.0003$), UPDRS 4 ($p = 0.0003$) PDQ-8 ($p = 0.001$) and NMSS total ($p = 0.0003$, Table 2). The LDE increased significantly during follow-up in the C group (1028.18 ± 388.76 to 1154.54 ± 407.09 mg, $p = 0.001$) while in the Apo, LDE decreased significantly (1077.81 ± 446.26 to 458.75 ± 282.29 , $p < 0.0001$) with improvement affecting more than 90% of patients and large effect size. In the Apo group, moderate to large effect size on the NMSS domains of sleep, mood/apathy, attention, gastrointestinal, urinary, sexual, and miscellany was evident (Table 3).

In the Apo group, there was a high correlation between change in UPDRS 3 and PDQ-8 ($r_s = 0.85$) and moderate between change in NMSS and PDQ-8

Table 1
Changes following Apomorphine infusion and continuing conventional therapy (comparator) in motor, non-motor, and quality of life dimensions

	Control			Apomorphine		
	Baseline	Follow-up	<i>p</i>	Baseline	Follow-up	<i>p</i>
UPDRS-Motor exam	20.06 (9.68)	19.35 (12.80)	0.69	36.94 (11.42)	15.35 (8.21)	0.0003
UPDRS-Complications	7.93 (5.43)	7.00 (4.46)	0.48	10.00 (6.43)	3.53 (3.52)	0.0003
NMSS-Cardiovascular	1.29 (2.97)	1.18 (2.90)	0.45	4.65 (5.63)	2.76 (3.51)	0.03
Sleep	12.29 (9.58)	12.06 (9.32)	0.90	22.06 (11.47)	10.71 (9.63)	0.0003
Mood/apathy	8.35 (10.33)	8.06 (8.78)	0.79	22.76 (19.85)	11.29 (13.04)	0.0005
Perceptual	2.23 (5.03)	2.59 (6.26)	0.90	4.59 (6.92)	1.88 (3.35)	0.04
Attention	6.00 (8.40)	7.18 (7.76)	0.16	12.82 (9.62)	8.71 (7.75)	0.006
Gastrointestinal	5.94 (5.97)	7.12 (6.49)	0.24	7.35 (7.35)	4.41 (5.11)	0.002
Urinary	4.29 (3.57)	6.23 (4.26)	0.06	10.70 (8.93)	5.71 (6.72)	0.001
Sexual	3.12 (6.58)	3.29 (6.12)	0.97	2.53 (5.96)	2.00 (3.94)	0.42
Miscellany	4.12 (5.67)	4.29 (5.55)	0.61	18.47 (14.54)	9.47 (9.70)	0.0003
NMSS-Total score	47.65 (43.40)	52.00 (37.65)	0.22	105.94 (65.43)	56.94 (45.39)	0.0003
PDQ-8	35.84 (23.10)	44.85 (17.57)	0.02	55.70 (19.80)	32.35 (21.54)	0.001

Benjamini-Hochberg correction: $p < 0.027$; UPDRS: Unified Parkinson's Disease Rating Scale; NMSS: Non-Motor Symptoms Scale; PDQ-8: Parkinson's Disease Questionnaire-8 items.

($r_s = 0.44$). The NNT was lower than 2 for >1 SEM improvement of UPDRS 3, NMSS, and PDQ-8 total scores. The average NNT for NMSS domains was 3.95 (Table 4).

DISCUSSION

Our key observations from this study are as follows:

1. There was a large beneficial effect of apomorphine infusion on the whole of NMSS observed along over one year average follow-up (Tables 2 and 3).

2. Patients with visual hallucinations, illusions and paranoid ideations reported no worsening after apomorphine infusion.
3. Apomorphine infusion resulted in significant improvement in some specific NMS such as hyperhidrosis, nocturia, urgency of micturition, and fatigue.

Our “real life” study included patients who may be excluded in randomised clinical trials and so we used a pragmatic post-treatment surveillance method. The follow-up period is therefore variable as clinical appointments were based on the discretion of the clinicians or local arrangement methods as also reported

Table 2
Change in the items of Non-Motor Symptoms Scale (Apomorphine infusion group)

Items & Domains	Baseline	Follow-up	<i>p</i> *
Cardiovascular			
1 Light-headedness	3.53 ± 3.98	2.29 ± 3.06	0.03
2 Fainting	1.23 ± 2.63	0.47 ± 0.94	0.27
Sleep/Fatigue			
3 Daytime sleeping	4.12 ± 4.44	2.88 ± 2.82	0.06
4 Fatigue	7.29 ± 3.93	2.82 ± 2.86	0.0004†
5 Difficult falling asleep	5.12 ± 5.04	2.47 ± 3.22	0.008†
6 Restless legs	5.53 ± 4.02	2.53 ± 3.54	0.003†
Mood/Cognition			
7 Lost interest surroundings	2.94 ± 4.25	0.82 ± 2.13	0.009†
8 Lack motivation	3.65 ± 4.26	1.18 ± 1.74	0.009†
9 Nervous	6.12 ± 5.43	3.23 ± 3.47	0.003†
10 Sad	5.53 ± 4.98	2.88 ± 3.71	0.001†
11 Flat mood	2.23 ± 2.93	1.47 ± 2.87	0.09
12 Difficult experiencing pleasure	2.76 ± 4.38	1.71 ± 3.22	0.015†
Perceptual problems			
13 Hallucinations	1.41 ± 2.45	0.53 ± 1.12	0.09
14 Delusions	1.53 ± 2.72	1.00 ± 1.87	0.3
15 Double vision	1.65 ± 3.26	0.35 ± 1.06	0.08

Table 2
(Continued)

	Items & Domains	Baseline	Follow-up	<i>p</i> *
	Cardiovascular			
	Attention/Memory			
16	Problems with concentration	5.47 ± 4.49	3.29 ± 3.69	0.002†
17	Forget recent events	4.06 ± 3.53	2.88 ± 2.59	0.1
18	Forget doing things	3.29 ± 3.51	2.53 ± 2.65	0.2
	Gastrointestinal			
19	Dribbling saliva	2.23 ± 2.36	1.35 ± 1.97	0.015†
20	Swallowing	2.00 ± 2.52	1.00 ± 1.58	0.026
21	Constipation	3.12 ± 4.03	2.06 ± 3.44	0.026
	Urinary			
22	Urgency	3.71 ± 3.88	1.88 ± 2.87	0.005†
23	Frequency	2.59 ± 3.00	1.41 ± 3.32	0.015†
24	Nocturia	4.41 ± 3.78	2.41 ± 2.67	0.005†
	Sexual function			
25	Altered interest in sex	1.59 ± 3.41	0.82 ± 1.74	0.05
26	Problems having sex	0.94 ± 3.01	1.18 ± 2.65	0.5
	Miscellaneous			
27	Unexplained pains	1.76 ± 3.99	2.59 ± 4.11	0.6
28	Lost taste/smell	4.41 ± 4.27	3.41 ± 4.18	0.05
29	Change in weight	3.00 ± 4.55	1.29 ± 2.23	0.05
30	Excessive sweating	7.59 ± 4.43	3.23 ± 3.77	0.001†

*Wilcoxon test; †Significant after Benjamini-Hochberg correction, *p* < 0.025.

from a large multicentre study of apomorphine infusion [8].

Apomorphine demonstrated benefit with a moderate to large effect size in all domains of NMSS (Table 3) along with UPDRS and PDQ-8. These data for the first time indicate efficacy of apomorphine infusion on NMS and improvement in HRQoL similar to the effect of intra-jejunal levodopa gel infusion in advanced PD as shown in a previous report from our group [28]. The NNT confirmed a beneficial effect for most patients,

both for motor and non-motor manifestations (Table 4).

We also report non-worsening of the perceptual and attention domain of NMSS (Tables 2 and 3) following apomorphine, contrary to what is to be expected with the use of a potent dopamine agonist in advanced PD. Apomorphine may have a psychotropic action based on its constituent anti-psychotic piperidine moiety [2, 3, 7, 29]. Improvement of psychosis was reported in 1978 in a placebo controlled study of apomorphine in 18 chronic schizophrenic patients [30]. In PD, open label

Table 3
Effect of the intervention (Apomorphine infusion group) in relation to motor, non motor and quality of life measures

	% of patients			Δ FU-B	Relative change	Effect Size	Standardized Response Mean
	I	NC	W				
UPDRS-Motor exam	100	0	0	-21.6 (9.56)	-58.44	1.89	2.26
UPDRS-Complications	94	6	0	-6.47 (4.91)	-64.70	1.01	1.32
NMSS-Cardiovascular	29	71	0	-1.88 (3.71)	-40.86	0.33	0.51
Sleep	100	0	0	-11.35 (6.93)	-51.45	0.99	1.64
Mood/apathy	82	18	0	-11.47 (10.90)	-50.39	0.58	1.05
Perceptual	41	53	6	-2.71 (5.19)	-59.04	0.39	0.52
Attention	71	23	6	-4.12 (7.31)	-32.14	0.43	0.56
Gastrointestinal	59	41	0	-2.94 (3.77)	-40.00	0.40	0.78
Urinary	71	29	0	-5.00 (5.26)	-46.70	0.56	0.95
Sexual	82	6	12	-0.53 (2.50)	-20.95	0.78	1.12
Miscellany	94	6	0	-9.00 (10.80)	-48.73	0.62	0.83
NMSS-Total score	100	0	0	-49.00 (36.62)	-46.25	0.75	1.34
PDQ-8	94	0	6	-23.34 (17.75)	-41.92	1.18	1.31

UPDRS: Unified Parkinson's Disease Rating Scale.

NMSS: Non-Motor Symptoms Scale; PDQ-8: Parkinson's Disease Questionnaire-8 items; I: Improved; NC: No changed; W: Worsened; Δ FU-B: Difference follow-up – baseline; mean (SD); For Effect size and Standardized Response Mean: 0.20–0.49, “small” effect; 0.50–0.79, “moderate” effect; and ≥0.80, “large” effect.

Table 4
NNT in patients treated with Apomorphine

	SEM	% of patients improving >1 SEM	NNT
UPDRS-Motor exam	10.48	82.35	1.21
UPDRS-Complications	5.34	47.06	2.12
NMSS- Cardiovascular	3.33	23.53	4.25
Sleep	8.11	58.52	1.71
Mood/apathy	11.74	41.18	2.43
Perceptual	4.89	17.65	5.67
Attention	6.08	29.41	3.40
Gastrointestinal	3.68	29.41	3.40
Urinary	5.21	35.29	2.83
Sexual	2.06	11.76	8.50
Miscellany	10.28	29.41	3.40
NMSS-Total score	42.40	52.94	1.90
PDQ-8	15.46	70.60	1.42

NNT = (1/% Improved) × 100; (n = 17).

studies had suggested safety of apomorphine infusion in those with neuropsychiatric problems on oral therapy [29, 31]. Tolerability of apomorphine infusion in our study may also be partly due to a significant reduction in LDE and cessation of therapy with other oral dopamine agonists, except the use of rotigotine patch in some cases.

The NMSS has been validated in over 600 patients and allows assessment of specific NMS within one instrument [13, 14]. Using NMSS, we report significant beneficial effects on sleep disturbances such as onset insomnia and restless legs while excessive daytime sleepiness was not worsened. Other NMS effects included a significant improvement in fatigue, motivation, anxiety, flat mood, anhedonia, attention deficit, dribbling of saliva, urinary dysfunction, particularly

urgency and nocturia, and hyperhidrosis (Table 2). Some of these NMS may have an underlying dopaminergic basis while in others the improvements indicate successful amelioration of subtypes of non-motor off periods [32]. Our data also supports data from previous open-label studies reporting a beneficial effect of apomorphine infusion on sleep (notably severe insomnia), anhedonia and night time pain and cramps suggestive of restless legs syndrome [11, 12, 29, 33].

This is not a randomised controlled study and we do not have a true control group. However, data from randomised controlled trials may occasionally have limited external validity [34]. We studied small number of patients although by comparison, the only other comparative study addressing apomorphine and deep brain stimulation included 13/12 patients only [35]. Similarly, the only comparative study of apomorphine versus levodopa infusion included 4 patients only [36]. In our study, a moderate to large effect of apomorphine was evident with the current sample size.

In conclusion, this observational post-treatment surveillance based study indicates the non-motor effects of apomorphine infusion. Specific non-motor effects of apomorphine are highlighted, calling for controlled studies with non-motor primary endpoints.

ACKNOWLEDGMENTS

This study forms a part of post treatment surveillance in real life study undertaken with EUROPAR, a non-profit academic group across Europe affiliated

Authors	Related to this article	Not related to this article
Pablo Martinez-Martin	None	GSK (2) Boehringer-Ingelheim, Novartis, Solvay, and Cephalon (5) Movement Disorder Society (6)
K. Ray Chaudhuri	None	GSK, Solvay, Boehringer-Ingelheim, Cephalon (2, 3, 5)
Regina Katzenschlager	None	UCB, Britannia, Teva (3, 5) UCB, Boehringer-Ingelheim (6).- GSK, UCB, TevaLundbeck, Santhera (2)
Per Odin	None	GSK, UBC, Boehringer, Cephalon, Abbott, Merck, Novartis (5) Boehringer – Ingelheim, Abbott, Cephalon, Ibsen, Orion – Pharma (3, 5)
Angelo Antonini	None	GSK, UCB, Boehringer – Ingelheim, Abbott, (2)
Tove Henriksen	None	Lundbeck Pharma (2), Abbott (2), Britannia (2), Orion – pharma (5)
Yogini Naidu, Susanne Tluk, Prashanth Reddy, Chandni Chandiramani	None	UCB, Abbott (3, 6)
Anne Martin	None	UCB, Abbott, GSK (3, 6)
Antonia Todorora	None	None

Funding category: 1. Stock Ownership in medically-related fields; 2. Consultancies; 3. Advisory Boards; 4. Partnerships; 5. Honoraria; 6. Grants; 7. Intellectual Property Rights; 8. Expert Testimony; 9. Employment; 10. Contracts; 11. Royalties; 12. Other.

with the European Parkinson's Disease Association. EUROPAR has received educational grants from Boehringer-Ingelheim, Britannia, UCB, Solvay and Parkinson's disease non-motor group.

AUTHOR'S ROLES

1. Research project: A. Conception, B. Organization, C. Execution;
2. Statistical Analysis: A. Design, B. Execution, C. Review and Critique;
3. Manuscript: A. Writing of the first draft, B. Review and Critique;

Pablo Martinez-Martin: 1A, 1B, 2A, 2B, 3.

A Antonini: 1B, C, 3 B

T Henriksen: 1C, 3B

R Katzenschlager: 1C, 3B

P Odin: 1C, 3B

Y Naidu: 1B, C, 3B

S Tluk: 1B, C, 3B

P Reddy: 1B, C, 3B

A Todorova: 1B, C, 3B

C Chandiramani: 1B, 3B

A Martin: 1B, C, 3B

K. Ray Chaudhuri: 1A, B, C, 2C, 3A B.

FINANCIAL DISCLOSURES

The authors report no conflicts of interest or competing financial interests with regards to this manuscript.

REFERENCES

- [1] Lees A, & Turner K (2002) Apomorphine for Parkinson's disease. *Practical Neurology*, **2**, 280-286.
- [2] Poewe W, & Wenning GK (2000) Apomorphine: An underutilized therapy for Parkinson's disease. *Mov Disord*, **15**(5), 789-794.
- [3] Chaudhuri KR, & Clough C (1998) Subcutaneous apomorphine for Parkinson's disease, Effective yet underused. *British Medical Journal*, **316**, 641.
- [4] National Institute for Health and Clinical Excellence. Clinical Guideline 35 Parkinson's disease: Diagnosis and management in primary and secondary care (2006) Available at: <http://www.nice.org.uk/nicemedia/pdf/cg035niceguideline.pdf> Accessed. 27 June, 2010.
- [5] Grosset DG, Macphie GJA, & Nairn M (2010) On behalf of the Guideline Development Group. Diagnosis and pharmacological management of Parkinson's disease: Summary of SIGN guidelines. *BMJ*, **340**, b5614.
- [6] Katzenschlager R, Hughes A, Evans A, Manson AJ, Hoffman M, & Swinn L et al. (2005) Continuous subcutaneous apomorphine therapy improves dyskinesias in Parkinson's disease: A prospective study using single-dose challenges. *Mov Disord*, **20**(2), 151-157.
- [7] Manson AJ, Turner K, & Lees AJ (2002) Apomorphine monotherapy in the treatment of refractory motor complications of Parkinson's disease: Long-term follow-up study of 64 patients. *Mov Disord*, **17**(6), 1235-1241.
- [8] Garcia Ruiz P, Ignacio SA, Pensando BA, Garcia AC, Frech FA, & Lopez MA et al. (2008) Efficacy of long term continuous subcutaneous apomorphine infusion in advanced Parkinson's disease with motor fluctuations: A multicenter study. *Mov Disord*, **23**(8), 1130-1136.
- [9] Stocchi F, Vacca L, De Pandis MF, Barbato L, Valente M, & Ruggieri S (2001) Subcutaneous continuous apomorphine infusion in fluctuating patients with Parkinson's disease: Long term results. *Neurol Sci*, **22**, 93-94.
- [10] Poewe W, Kleedorfer B, Wagner M, Bosch S, & Schelosky L (1993) Continuous subcutaneous apomorphine infusions for fluctuating Parkinson's disease. Long term follow up in 18 patients. *Adv Neurol*, **60**, 656-659.
- [11] Martinez-Martin P, Todorova A, Odin P, Henriksen T, Naidu Y, Tluk et al. (2010) The tolerability of apomorphine infusion in advanced Parkinson's patients with hallucinations and gastrointestinal non motor symptoms: Implications for treatment. A EUROPAR study. *Mov Disord*, **25**(Suppl 2), S421.
- [12] Naidu Y, Logishetty K, Martin A, Tluk S, Reddy P, Martinez-Martin P, Odin P, Henriksen T, & Chaudhuri KR (2009) Efficacy of apomorphine infusion on mood and depression aspects of non-motor symptoms in advanced Parkinson's disease? *Park Rel Disord*, **15**(Suppl 2), S130 (P2.149).
- [13] Chaudhuri KR, Martinez-Martin P, Brown RG, Sethi K, Stocchi F, & Odin P et al. (2007) The metric properties of a novel non-motor symptoms scale for Parkinson's disease: Results from an international pilot study. *Mov Disord*, **22**, 1901-1911.
- [14] Martinez-Martin P, Rodriguez-Blazquez C, Abe K, Bhatlacharya KB, Bloem BR, & Carod-Artal FJ et al. (2009) International study on the psychometric attributes of the non-motor symptoms scale in Parkinson disease. *Neurology*, **73**, 1584-1591.
- [15] Hughes AJ, Daniel SE, Kilford L, & Lees AJ (1992) Accuracy of clinical diagnosis of idiopathic Parkinson's disease: A clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry*, **55**, 181-184.
- [16] Fahn S, & Elton RL (1987) Members of the UPDRS Development Committee. Unified Parkinson's disease Rating Scale. En: Fahn S, Marsden CD, Calne DB, Goldstein M, eds. Recent developments in Parkinson's disease, Macmillan Healthcare Information, Florham Park, NJ. **2**, 153-164.
- [17] Hoehn MM, & Yahr MD (1967) Parkinsonism: Onset, progression, and mortality. *Neurology*, **17**, 427-442.
- [18] Jenkinson C, Fitzpatrick R, Peto V, Greenhall R, & Hyman N (1997) The PDQ-8: Development and validation of a short-form Parkinson's disease questionnaire. *Psychol Health*, **12**, 805-814.
- [19] Benjamini Y, & Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J Roy Statistical Society*, **57**, 289-300.
- [20] Swinscow TDV (1997) Correlation and Regression, BMJ Publishing Group, Statistics at square one, 9th edition. University of Southampton.
- [21] Kazis L, Anderson J, & Meenan R (1989) Effect sizes for interpreting changes in health status. *Med Care*, **27**(Suppl), 178-189.
- [22] Husted JA, Cook RJ, Farewell WT, & Gladman DD (2000) Methods for assessing responsiveness: A critical review and recommendations. *J Clin Epidemiol*, **53**, 459-468.

- [23] Crosby RD, Kolotkin RL, & Williams GR (2003) Defining clinically meaningful change in health-related quality of life. *J Clin Epidemiol*, **56**, 395-407.
- [24] Guyatt GH, Osoba D, Wu AW, Wyrwich K, Norman G (2002) The Clinical Significance Consensus Meeting Group Methods to explain the clinical significance of health status measures. *Mayo Clin Pro*, **77**, 371-383.
- [25] Schünemann HJ, Akl EA, & Guyatt GH (2006) Interpreting the results of patients reported outcome measures in clinical trials: The clinician's perspective. *Health Qual Life Outc*, **4**, 62.
- [26] Wyrwich KW, Bullinger M, & Aaronson N et al. (2005) Estimating clinically significant differences in quality of life outcomes. *Qual Life Res*, **14**, 285-295.
- [27] Rejas J, Pardo A, & Ruiz MA (2008) Standard error of measurement as a valid alternative to minimally important difference for evaluating the magnitude of changes in patient-reported outcomes measures. *J Clin Epidemiol*, **61**, 350-356.
- [28] Honig H, Antonini A, Martinez-Martin P, Forgacs I, Faye GC, & Fox T et al. (2009) Intrajejunal levodopa infusion in Parkinson's disease: A pilot multicenter study of effects on nonmotor symptoms and quality of life. *Mov Disord*, **24**(0), 1468-1474.
- [29] van Laar T, Postma AC, & Drent M (2010) Continuous subcutaneous infusion of Apomorphine can be used safely in patients with Parkinson's disease and pre-existing visual hallucinations. *Parkinsonism Relat Disord*, **16**, 71-72.
- [30] Tamminga CA, Schaffer MH, Smith RC, & Davis JM (1978) Schizophrenic symptoms improve with apomorphine. *Science*, **200**, 567-568.
- [31] Ellis C, Lemmens G, Parkes JD, Abbott RJ, Pye IF, & Leigh PN et al. (1997) Use of apomorphine in parkinsonian patients with neuropsychiatric complications to oral treatment. *Parkinsonism Relat Disord*, **3**(2), 103-107.
- [32] Witjas T, Kaphan E, Azulay JP, Blin O, Ceccaldi M, Pouget J, Poncet M, & Chérif AA (2002) Nonmotor fluctuations in Parkinson's disease: Frequent and disabling. *Neurology*, **59**(3), 408-413.
- [33] Reuter I, Ellis CM, & Chaudhuri KR (1999) Nocturnal subcutaneous apomorphine infusion in Parkinson's disease and restless legs syndrome. *Acta Neurol Scand*, **100**, 163-167.
- [34] Relton C, Togerson D, O'Cathain AO, & Nicholl J (2010) Rethinking pragmatic randomised controlled trials: Introducing the cohort multiple randomised controlled trial design. *BMJ*, **340**, 963-967.
- [35] De Gaspari D, Siri C, Landi A, Cilia R, Bonetti A, & Natuzzi F et al. (2006) Clinical and neuropsychological follow up at 12 months in patients with complicated Parkinson's disease treated with subcutaneous apomorphine infusion or deep brain stimulation of the subthalamic nucleus. *J Neurol Neurosurg Psychiatry*, **77**(4), 450-453.
- [36] Nyholm D (2009) Comparison of apomorphine and levodopa infusions in four patients with Parkinson's disease with symptom fluctuations. *Acta Neurol Scand*, **119**(5), 345-348.